

Short report

Hedonic and disgust taste perception in borderline personality disorder and depression

Gonzalo Arrondo, Graham K. Murray, Emma Hill, Bence Szalma, Krishna Yathiraj, Chess Denman and Robert B. Dudas

Summary

Depression and borderline personality disorder (BPD) are both thought to be accompanied by alterations in the subjective experience of environmental rewards. We evaluated responses in women to sweet, bitter and neutral tastes (juice, quinine and water): 29 with depression, 17 with BPD and 27 healthy controls. The BPD group gave lower pleasantness and higher disgust ratings for quinine and juice compared with the control group; the depression group did not differ significantly from the control group. Juice disgust ratings were related to self-disgust in BPD, suggesting close

links between abnormal sensory processing and self-identity in BPD.

Declaration of interest

None.

Copyright and usage

© The Royal College of Psychiatrists 2015. This is an open access article distributed under the terms of the Creative Commons Attribution (CC BY) licence.

Reduced enjoyment of pleasant experiences and increased impact of negative emotions are observed clinically in depression and in borderline personality disorder (BPD).¹ Surprisingly, previous laboratory studies do not support a link between depression and reduced pleasantness ratings to biological rewards, such as sucrose solutions (for example see Swiecicki *et al*²). However, patients with depression could be more sensitive to negative primary inputs.³ Few studies have used aversive stimuli in the evaluation of taste in depression, and no studies have been carried out to evaluate hedonic ratings of tastes in BPD in spite of clinically observable aberrant emotional processing and increased state and trait disgust.⁴ We hypothesised that patients with BPD and depression would differ from healthy controls in their pleasantness and disgust ratings to positive and especially negative taste stimuli.

Method

A total of 29 women with DSM-IV⁵ major depressive disorder, 17 women with DSM-IV BPD and 27 female healthy controls took part in the study, which was approved by the Cambridgeshire 4 National Health Service research ethics committee; all participants provided written informed consent. Additional details, including statistical data, are provided in the online supplement. Evaluation of taste consisted of participants taking a sip, but not swallowing, from a cup with 10 ml of orange juice, quinine dihydrochloride at 0.006 mol/L or water. Participants had to maintain the liquid in the mouth for 5 s, rate the disgust and pleasantness produced using two visual scales (online Fig. DS1) and rinse their mouths with water. Order of liquids was counterbalanced across participants. Clinical evaluation was completed prior to the taste experiment. Statistical analysis (see online supplement for details) aimed to evaluate the association between taste disgust and disgust as measured using two clinical rating scales: the Self-Disgust Scale (SDS)⁶ and the Disgust Scale Revised (DSR).⁷

Results

Overall differences between the three conditions in pleasantness and disgust ratings followed predictions across all participants. Quinine was highly unpleasant and disgusting, juice was highly pleasant and not disgusting and water was neither pleasant nor disgusting. Pleasantness and disgust ratings correlations were all significant. Regarding differences between groups, the BPD group

rated both quinine and juice (but not water) as more unpleasant and disgusting than the control group, but no differences were found between the depression and control groups (all $P < 0.05$; see online Tables DS8 and DS9, and Fig. 1). Increased self-disgust was significantly correlated (Spearman's $\rho = 0.5$) with greater disgust ratings after the intake of orange juice in the BPD group (Table DS10). However, disgust propensity did not correlate with ratings in this group and all correlations were non-significant in the depression group.

Discussion

We found that in the BPD group there were abnormal pleasantness and disgust ratings after the intake of biological stimuli, whereas no differences between the depression and control groups were found. Our findings indicate that the hedonic experience of both positive and negative taste stimuli is negatively biased in BPD. This novel result is in line with clinical findings in the disorder, as people with BPD report more dysphoric and less positive affective cognitive states.¹ Current diagnostic criteria for the disorder include affective instability, recurrent self-threatening behaviours and chronic feelings of emptiness, all of which could be related to negative perceptions of the environment.

The lack of evidence for a differential effect in the case of depression is also in line with most of the existing literature on enjoyment of pleasant tastes;² however, our study also shows that there were no differences between the depression group and the control group in evaluation of a disgusting taste. A limitation of our study and the prior depression studies is that sample sizes were small; hence, either there is no true difference in the ratings for chemosensory stimuli between people with depression and controls or the effect is small, suggesting that the basis of anhedonia reported in depression is complex. For example, it could be that clinically observed anhedonia in depression is primarily related to social anhedonia. Alternatively, clinical assessments of anhedonia may confound motivational, anticipatory and mnemonic aspects of enjoyment with consummatory 'in the moment' pleasure; the latter is assayed by our laboratory taste task and may be comparatively intact in depression.⁸

In the BPD group, questionnaire-measured self-disgust, but not disgust propensity, correlated with laboratory-rated disgust




Fig. 1 Scatter plot with the (a) juice and (b) quinine disgust ratings stratified by group.

The horizontal brackets with asterisks indicate significant group differences and the horizontal lines group medians. BPD, borderline personality disorder; MDD, major depressive disorder.

to juice stimuli. Self-disgust indicates a context-free negative evaluation of the self (shame feelings) and also negative views about one's actions (guilt),⁶ and it is greatly enhanced in BPD. Self-disgust may be felt as an embodied experience instead of an abstract sensation, to the point of producing negative physical sensations such as nausea, and is often triggered by external events.⁹ Our finding of an association between self-disgust and juice-disgust indicates close links between sensory processing and self-identity in BPD, and may suggest that basic physiological disturbances play a role in the origins of self-disgust in this disorder. Previous research indicates that self-disgust is correlated with overall symptom severity in BPD and eating disorders,¹⁰ which could indicate similar mechanisms within the two disorders.¹¹ We speculate that in BPD self-disgust is so heightened that it may impair the enjoyment of stimuli that are ordinarily considered as pleasant. Alternatively, a fundamental abnormality in processing external sensory stimuli may contribute to a negative sense of self in BPD. The less positive ratings could also be related to other group-specific factors, such as an increased history of trauma in the BPD group.^{4,12} Our results emphasise the significance of disgust – both of the self and of external stimuli – in BPD, and highlight a role for assessment of disgust in the diagnosis and management of this condition.⁴

Gonzalo Arrondo, PhD, Department of Psychiatry, University of Cambridge, Cambridge; **Graham K. Murray**, MD, PhD, MRCPsych, Department of Psychiatry, University of Cambridge, Cambridgeshire and Peterborough NHS Foundation Trust and Behavioural and Clinical Neuroscience Institute, University of Cambridge, Cambridge; **Emma Hill**, PhD, Cambridgeshire and Peterborough NHS Foundation Trust, Cambridge; **Bence Szalma**, BSc, Department of Psychiatry, University of Cambridge, Cambridge; **Krishna Yathiraj**, MRCPsych, Department of Psychiatry, University of Cambridge, Cambridge and Norfolk and Suffolk NHS Foundation Trust, Norwich; **Chess Denman**, MRCPsych, Cambridgeshire and Peterborough NHS Foundation Trust, Cambridge; **Robert B. Dudas**, MD, PhD, MRCPsych, Department of Psychiatry, University of Cambridge, Cambridgeshire and Peterborough NHS Foundation Trust, Behavioural and Clinical Neuroscience Institute, University of Cambridge, Cambridge, and Psychiatric Liaison Service, Ipswich Hospital, Norfolk and Suffolk NHS Foundation Trust, Norwich, UK

Correspondence: Graham K. Murray, Department of Psychiatry, University of Cambridge, Herchel Smith Building, Forvie Site, Robinson Way, Cambridge Biomedical Campus, Cambridge CB2 0SZ, UK. Email: gm285@cam.ac.uk

First received 20 Apr 2014, final revision 30 Oct 2014, accepted 7 Nov 2014

Funding

Supported by the Wellcome Trust (093875/Z/10Z and 09814/Z/11), Medical Research Council (G0701911 and G1000183), Isaac Newton Trust, Cambridgeshire and Peterborough NHS Foundation Trust, Talisman Trust, University of Cambridge and the NIHR Cambridge Biomedical Research Centre.

References

- 1 Reed LI, Fitzmaurice G, Zanarini MC. The course of positive affective and cognitive states in borderline personality disorder: a 10-year follow-up study. *Personal Ment Health* 2012; **6**: 281–91.
- 2 Swiecicki L, Zatorski P, Bzinkowska D, Sienkiewicz-Jarosz H, Szyndler J, Scinska A. Gustatory and olfactory function in patients with unipolar and bipolar depression. *Prog Neuropsychopharmacol Biol Psychiatry* 2009; **33**: 827–34.
- 3 Leppanen JM. Emotional information processing in mood disorders: a review of behavioral and neuroimaging findings. *Curr Opin Psychiatry* 2006; **19**: 34–9.
- 4 Schienle A, Haas-Krammer A, Schoggl H, Kapfhammer HP, Ille R. Altered state and trait disgust in borderline personality disorder. *J Nerv Ment Dis* 2013; **201**: 105–8.
- 5 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (4th edn) (DSM-IV)*. APA, 1994.
- 6 Overton PG, Markland FE, Taggart HS, Bagshaw GL, Simpson J. Self-disgust mediates the relationship between dysfunctional cognitions and depressive symptomatology. *Emotion* 2008; **8**: 379–85.
- 7 Olatunji BO, Williams NL, Tolin DF, Abramowitz JS, Sawchuk CN, Lohr JM, et al. The Disgust Scale: item analysis, factor structure, and suggestions for refinement. *Psychol Assess* 2007; **19**: 281–97.
- 8 Dillon DG, Rosso IM, Pechtel P, Killgore WD, Rauch SL, Pizzagalli DA. Peril and pleasure: an rdoc-inspired examination of threat responses and reward processing in anxiety and depression. *Depress Anxiety* 2014; **31**: 233–49.
- 9 Powell PA, Overton PG, Simpson J. The revolting self: an interpretative phenomenological analysis of the experience of self-disgust in females with depressive symptoms. *J Clin Psychol* 2014; **70**: 562–78.
- 10 Ille R, Schoggl H, Kapfhammer HP, Arendasy M, Sommer M, Schienle A. Self-disgust in mental disorders – symptom-related or disorder-specific? *Compr Psychiatry* 2014; **55**: 938–43.
- 11 Zanarini MC, Reichman CA, Frankenburg FR, Reich DB, Fitzmaurice G. The course of eating disorders in patients with borderline personality disorder: a 10-year follow-up study. *Int J Eat Disord* 2010; **43**: 226–32.
- 12 Rusch N, Schulz D, Valerius G, Steil R, Bohus M, Schmahl C. Disgust and implicit self-concept in women with borderline personality disorder and posttraumatic stress disorder. *Eur Arch Psychiatry Clin Neurosci* 2011; **261**: 369–76.



Data supplement to Arrondo et al. Hedonic and disgust taste perception in borderline personality disorder and depression. *Br J Psychiatry* doi: 10.1192/bjp.bp.114.150433

Online supplement

Previous literature on taste and major depressive disorder (MDD)

To the best of our knowledge seven previous published articles have studied if there are differences in the appraisal and/or recognition of taste stimuli between patients with depression and controls. We have included the main data from them in Table DS1.

The hypothesis behind these studies, namely that depression patients may perceive positive stimuli differently from the average population, is not new and studies date back as far as 1969. The number of participants has been small with clinical samples between 12 and 36 patients and control samples between 15 and 30 participants. The stimuli most typically evaluated are sucrose solutions (i.e. sweetness) and the sucrose concentrations have varied within a study, whereas specific works additionally studied the responses to bitter, sour, salty or citric stimuli. Regarding the measures obtained, most researchers have typically used either pleasantness or intensity ratings or threshold identification. The latter consists in participants indicating at which concentration they identify the presence of sucrose, hence obtaining an identification threshold in a typical psychophysical paradigm. The former measures are usually obtained by asking participants to indicate the pleasantness produced by the intake of the stimuli in a visual scale, or conversely, its intensity.

The first study in this manner evaluated the taste thresholds for different flavours in hospitalized patients and found that the recognition threshold in depression was heightened for all kinds of stimuli when compared to other psychiatric control patients. Moreover, they also found that the clinical symptoms correlated with this increased threshold and that recovery led to improvements in taste.¹³ A study nearly 20 years later found increased intensity thresholds, but surprisingly increased pleasantness ratings in depression group when compared to normal controls.¹⁴ Nevertheless this difference was not significant when only unipolar depression patients were considered. The subsequent studies were not been able to find differences in pleasantness ratings to positive stimuli between patients and controls.

See Table DS1 for a list of main relevant articles and a comparison of their most critical features.

Table DS1 Main relevant articles

| Authorship | Year | Depression group, <i>n</i> | Health control group, <i>n</i> | Measure | Flavours |
|-------------------------------|------|----------------------------|--------------------------------|---|-----------------------------------|
| Dichter et al ¹⁸ | 2010 | 12 | 15 | Pleasantness rating Intensity rating | Sweet |
| Swiecicki et al ² | 2009 | 21 | 30 | Pleasantness rating Threshold identification Flavour identification | Sweet Bitter Citric Sour |
| Berlin et al ¹⁵ | 1998 | 20 | 20 | Pleasantness rating Threshold identification | Sweet |
| Potts et al ¹⁷ | 1997 | ~* | ~* | Threshold identification | Sweet |
| Steiner et al ¹⁶ | 1993 | 21 | 16 | Pleasantness rating | ** |
| Amsterdam et al ¹⁴ | 1987 | 19 | 36 | Pleasantness rating Intensity rating | Sweet |
| Steiner et al ¹³ | 1969 | 21 | - | Threshold identification | Sweet Sour Salty Bitter |

If the studies included other clinical samples these are not detailed in the table.

Measure is the type of measure carried out, whereas flavour is the different flavours used in the study.

* Samples were changed during the study

** Authors have not been able to access the original article

Participants

We used structured diagnostic interview schedules, the MINI (Mini-International Neuropsychiatric Interview) and SCID-II (Structured Clinical Interview for DSM-IV, part II) to establish the diagnoses required for inclusion and exclude volunteers who did not meet criteria. BPD patients were outpatients recruited via the Complex Cases Service (CCS), a specialized personality disorders unit; participants with unipolar depression by newspaper advertisements, and healthy participants from the Medical Research Council Cognition and Brain Sciences Unit healthy volunteer panel and also via advertisements. All participants were interviewed by a psychiatrist (the last author of this paper) with expertise in personality disorders. BPD patients with current or past history of any formally diagnosed psychotic illness or current major depressive disorder, or dependence on a psychoactive substance, as per the MINI, were excluded. The presence of depressive symptoms (as opposed to a full-blown, co-morbid major depressive illness) did not lead to exclusion. The presence of other personality disorder traits, but not that of the full-blown disorder, was permitted. In the MDD group, any comorbid psychiatric conditions as per the MINI or SCID led to exclusion, but the presence of personality disorder traits, without the full-blown disorder, was permitted. In healthy volunteers, any history or presence of psychiatric or neurological illness led to exclusion. No participant had any history of epilepsy, serious head injury, serious medical conditions, physical problems requiring hospitalisation, or surgery in general anaesthesia in the previous 6 months. Furthermore, all participants were tested during the follicular phase of their menstrual cycle (days 3-10) to eliminate the potential confounding factor of differential emotional responding due to hormonal differences.

Ethics statement: A local NHS research ethics committee approved this research (Cambridgeshire 4 Research Ethics Committee, NHS National Research Ethics Service, reference number: 09/H0305/10). Written informed consent was obtained from each participant.

The age of the different groups was 31.8 (SD 7.8) for the healthy controls, 35.3 (SD 7.8) for the BPD and 35.6 (SD 8.8) for the MDD group (mean and standard deviations are provided). When compared to the controls the MDD group was significantly different ($t=-2.13$ $p=0.038$) while BPD patients' age was also close to being significantly different compared to the health control (HC) group ($t=-1.87$ $p=0.069$). There were no differences between patient groups ($t=-0.91$ $p=0.928$). Nevertheless it must be noted that whether statistically significant or not, these differences do not seem to be of real clinical relevance as the overall range of ages was not wide. Moreover any differences found between the HC and only one of the patients' groups are unlikely to be explained by the age, since both patient groups had an older mean age than controls. This is especially so in the case that differences were found only in the BPD group (as is our case), in which the statistical significance of the differences was smaller than the in MDD group.

Participants were advised not to have any coffee, tea, or "energy drinks" such as "*Red Bull*" during the 2 hours previous to the evaluation and not to smoke in the previous hour. They were questioned regarding the fulfilment of this requirement prior to the experiment. Additionally participants were questioned to ensure that none had taken any psychoactive substances in the previous 24 hours.

On the choice of tastes and visual scales

Orange juice was chosen as a positive stimulus, quinine as a negative one and water as a neutral condition. While previous studies have typically used sucrose solutions for the evaluation of the pleasantness of positive stimuli, we decided to include orange juice as a more typical positive stimulus in everyday life.

Although we expected a high (inverse) correlation between the disgust and pleasantness ratings the rationale for the inclusion of two separate measures is that it was hypothesized that the former might be more sensitive to find differences in taste in the BPD group as the disorder is characterized by the alteration of a broad spectrum of disgust processes.

Taste evaluation

Studies lasted for approximately half an hour, with the questionnaire phase lasting 20 minutes and the taste evaluation 10 more minutes.

Stimuli were quinine dihydrochloride, orange juice and water. Concentration of quinine dihydrochloride was 0.006 mol/L solution and was prepared in the local pharmacy at a higher concentration and further diluted following the pharmacist's instructions at the research site for the experiment. Orange juice was obtained from a common brand of orange squash following the manufacturer's recommendation regarding dilution with water. Tap water was used for the quinine and juice dilutions as well as for the water flavour.

Ten mls from each stimulus liquid were put into closed blank plastic disposable cups for every participant. Cups were numbered for ease of randomization.

Evaluation of taste consisted in volunteers taking a sip, but not swallowing from a cup with 10 ml of orange juice (J), quinine dihydrochloride at 0.006 mol/L (Q) or water (W). After putting the liquid in the mouth they had to maintain it there for 5 seconds (s) and then swallow or spit it out at their discretion. 30s later participants rated with a pen in two paper visual scales the pleasantness (20cm long, going from -10-very unpleasant- to 10 – very pleasant-) and disgust (10cm long, going from 0-not disgusting at all-, to 10-extremely disgusting) and finally rinsed their mouths with water after further 30s. There was at least a further 60s between flavour evaluations. Order of liquids was counterbalanced across subjects whereas clinical evaluation was completed prior to the taste experiment.

Prior to the taste evaluation participants held a clinical interview which included the Mini-international neuropsychiatric interview (MINI),¹⁹ and completed several clinical measures and questionnaires.

Statistical analysis

Non-parametric tests were used throughout because of the ceiling effect of the visual scales and hence the non-normality of the obtained data. We used a Friedman ANOVA for repeated measures to evaluate whether there was an effect of condition (i.e. beverage) in the pleasantness and disgust ratings; we compared the three groups with a Kruskal-Wallis analysis and, as we were especially interested in differences between the two clinical groups and the control group these were compared through planned Mann-Whitney *U*-tests. Rho Spearman correlations were used to evaluate the association between taste disgust and disgust as rated by the two clinical rating scales, the SDS⁶ and the DSR.⁷ Statistical analyses were conducted with

SPSS21 (IBM; Armonk, NY, US) running on a Dell Optiplex789 with Windows XP. The threshold for significance was set at $p < 0.05$ and tests were two-sided.

History of traumatization

There is some evidence linking history of psychological traumatization and abnormal disgust ratings both regarding self-disgust and also food or fluid-related disgust.^{4,12} Since history of psychological trauma and post-traumatic stress disorder is more commonly seen in BPD patients than in the normal population it could be mediating the differences between groups. Subjects were divided into participants which had a history of trauma and those who did not, depending on answers to questions H1 and H2 (PTSD section) of the MINI questionnaire (H1: Have you ever experienced or witnessed or had to deal with an extremely traumatic event that included actual or threatened death or serious injury to you or someone else?; H2: Did you respond with intense fear, helplessness or horror?; participants answering affirmatively to both questions were considered to have a history of trauma). The number of participants with trauma history was 1, 0, and 9 for the HC, MDD and BPD group respectively. We then compared disgust and pleasantness ratings, as well as disgust scales, between the trauma and no trauma subgroups of the BPD participants in an exploratory analysis to see if trauma history could be influencing our results (See Table DS12). There were no differences in the disgust questionnaires between subgroups and similarly we did not find differences in the disgust ratings for quinine and juice, which were the measurements that differentiated BPD participants. On the other hand, BPD participants with traumatization history had increased disgust ratings for water and reduced pleasantness ratings when tasting quinine. Therefore, while history of trauma does not seem to be directly involved in the results found in this study, it is an important construct that should be controlled and evaluated in future studies,^{4,11} ideally by use of a quantitative psychometric instrument.

Table DS2 Psychiatric medication

| | BPD (N=17) | MDD (N=29) |
|--------------------------|-------------------|-------------------|
| Any medication | 13 (76.5) | 19 (65.5) |
| Any antidepressant | 8 (47.1) | 17 (58.6) |
| SSRI | 5 (29.4) | 15 (51.7) |
| Non SSRI antidepressants | 4 (23.5) | 4 (13.8) |
| Antipsychotic | 3 (17.6) | 0 (0) |
| Mood stabilizers | 3 (17.6) | 0 (0) |
| Benzodiazepines | 5 (29.4) | 0 (0) |

BPD is Borderline Personality Disorder and MDD is Major Depressive Disorder. Number of patients in each clinical group (with percentage in brackets) taking a given class of medications. Healthy controls did not take any psychiatric medication per inclusion criteria.

Table DS3 Clinical scales

| | HC | BPD | MDD |
|------|------------------|--------------|-----------------|
| SDS | 21 (16.25-25) | 61 (52.5-69) | 42 (37.5-50.5) |
| DSR | 10.5 (6.75-12.5) | 15 (13-20) | 13 (8.5-16) |
| BDI | 1 (0-3) | 24 (9-34.5) | 19 (13-26) |
| HRSD | 0 (0-1) | 15 (5-16) | 21.5 (18.75-27) |

SDS is Self-Disgust Scale,⁶ DSR is Disgust Scale Revised,^{7,20} BDI is the Beck Depression Inventory-II;²¹ and HRSD is the Hamilton Rating Scale for Depression Scale.²² Median with interquartile ranges (in brackets) are given. HC is Healthy Controls, BPD Borderline Personality Disorder and MDD is Major Depressive Disorder.

Table DS4 Correlation between pleasantness and disgust

| | Juice | Water | Quinine |
|---------|--------|--------|---------|
| Rho | -0.261 | -0.434 | -0.803 |
| p-value | 0.026 | <0.001 | <0.001 |

Spearman correlations between pleasantness and disgust scales in the whole group of participants.

Table DS5 Effect of condition on pleasantness (Friedmann ANOVA)

| | TS | SE | Std TS | p-value | Adj. p-value |
|------------------------|---------|-------|--------|---------|--------------|
| Friedmann Anova | 116.561 | - | - | <0.001 | - |
| J vs W | 0.664 | 0.166 | 4.014 | <0.001 | <0.001 |
| Q vs W | -1.096 | 0.166 | 6.621 | <0.001 | <0.001 |
| Q vs J | -1.760 | 0.166 | 10.635 | <0.001 | <0.001 |

TS is the Test Statistic (Chi-Square for Friedman Anova, and W Wilcoxon for the post-hoc comparisons), SE is standard Error, Std TS is the standardized Test Statistic, Adj p-value is the adjusted significance for multiple comparisons (Dunn-Bonferroni procedure²³). J is juice, W is water, and Q is quinine.

Table DS6 Effect of condition on disgust (Friedman ANOVA)

| | TS | SE | Std TS | p-value | Adj. p-value |
|------------------------|--------|-------|--------|---------|--------------|
| Friedmann Anova | 111.85 | - | - | <0.001 | - |
| J vs W | -0.116 | 0.166 | -0.703 | -0.482 | 1 |
| Q vs W | 1.349 | 0.166 | -8.152 | <0.001 | <0.001 |
| Q vs J | -1.466 | 0.166 | -8.855 | <0.001 | <0.001 |

TS is the Test Statistic (Chi-Square for Friedman Anova, and W Wilcoxon for the post-hoc comparisons), SE is standard Error, Std TS is the standardized Test Statistic, Adj p-value is the adjusted significance for multiple comparisons (Dunn-Bonferroni procedure²³). J is juice, W is water, and Q is quinine.

Table DS7 Effect of group on ratings (Kruskal-Wallis test)

| Liquid | Measure | TS | p-value |
|---------|--------------|------|---------|
| Juice | Pleasantness | 5.30 | 0.070 |
| | Disgust | 5.24 | 0.073 |
| Water | Pleasantness | 2.62 | 0.269 |
| | Disgust | 0.66 | 0.719 |
| Quinine | Pleasantness | 6.53 | 0.038 |
| | Disgust | 8.26 | 0.016 |

TS is the Test Statistic (Kruskal-Wallis H)

Table DS8 Effect of group on pleasantness (Mann-Whitney U tests)

| | WG | HC | BPD | MDD | BPD vs HC | MDD vs HC |
|---------|----------------|--------------------|-------------------|--------------------|---------------------------------|---------------------------------|
| Juice | 4 (1.5 — 6.1) | 4.3(3.1—6.9) | 2.1 (-0.6 — 5) | 3.8 (1.4 — 5.9) | 135 (SE=41.47; p=0.023) | 311.5 (SE=60.97; p=0.189) |
| Water | 0 (-0.1 —0.1) | 0 (-0.4 — 0) | 0 (-1 — 0) | 0 (0 —1.4) | 212 (SE=38.81; p=0.652) | 466 (SE=57.85; p=0.198) |
| Quinine | -8 (-9.6 — -6) | -7.1 (-8.2 — -5.4) | -9.5 (-10 — -7.4) | -7.9 (-9.5 — -6.1) | 129.5 (SE=41.29; p=0.015) | 326.5 (SE=60.90; p=0.286) |

WG is Whole Group, HC Healthy Controls, BPD Borderline Personality Disorder and MDD is Major Depressive Disorder. In 4 first columns median with interquartile ranges (in brackets) are given. Last 2 columns are U Mann Whitney results, test standard error (SE) and its associated p value.

Table DS9 Effect of group on disgust (Mann-Whitney U tests)

| | WG | HC | BPD | MDD | BPD vs HC | MDD vs HC |
|----------------|---------------|-------------|-------------|---------------|---------------------------------|----------------------------|
| Juice | 0(0—0.1) | 0 (0—0) | 0 (0—1.4) | 0 (0—0.1) | 305.5 (SE=34.28; p=0.027) | 434 (SE=47.50; p=0.371) |
| Water | 0 (0—0.5) | 0 (0—0.6) | 0.1 (0—1.3) | 0 (0—0.45) | 253 (SE=38.4; p=0.612) | 380 (SE=55.55; p=0.836) |
| Quinine | 7.5 (4.1-9.3) | 6.2 (3—8.1) | 9 (6.9—10) | 7.4 (4.7—9.3) | 342 (SE=41.4; p=0.007) | 491 (SE=60.95; p=0.103) |

WG is Whole Group, HC Healthy Controls, BPD Borderline Personality Disorder and MDD is Major Depressive Disorder. In 4 first columns median with interquartile ranges (in brackets) are given. Last 2 columns are U Mann Whitney results, test standard error (SE) and its associated p value.

Table DS10 Correlations between disgust scales and disgust ratings in BPD (Spearman)

| | Juice | | Water | | Quinine | |
|------------|-------|---------|--------|---------|---------|---------|
| | Rho | p-value | Rho | p-value | Rho | p-value |
| SDS | 0.501 | 0.037 | 0.111 | 0.670 | 0.295 | 0.251 |
| DSR | 0.151 | 0.562 | -0.101 | 0.701 | -0.006 | 0.982 |

SDS is Self-Disgust Scale⁶, DSR is Disgust Scale Revised.^{7,20} J is juice, W is water and Q is quinine. Rho is the Spearman correlation.

Table DS11 Correlations between disgust scales and disgust ratings in MDD (Spearman)

| | Juice | | Water | | Quinine | |
|------------|-------|---------|--------|---------|---------|---------|
| | Rho | p-value | Rho | p-value | Rho | p-value |
| SDS | 0.131 | 0.498 | -0.104 | 0.590 | -0.003 | 0.988 |
| DSR | 0.112 | 0.563 | -0.262 | 0.170 | -0.181 | 0.347 |

SDS is Self-Disgust Scale,⁶ DSR is Disgust Scale Revised^{7,20} J is juice, W is water and Q is quinine. Rho is the Spearman correlation.

Table DS12 Effects of traumatization in the BPD group(Mann-Whitney U tests)

| | No trauma | Trauma | Trauma vs. No trauma |
|-----------------------------|-------------------|------------------|------------------------------|
| Juice pleasantness | 2.5 (-1.2 – 8.2) | 1.5 (-2.9 – 5.0) | 26.5 (SE = 9.43; p = 0.606) |
| Juice disgust | 0.6 (0.0 – 2.0) | 0.0 (0.0 – 2.7) | 24.5 (SE = 8.84; p = 0.470) |
| Water pleasantness | -0.8 (-5.2 – 0.0) | 0.0 (0.0 - 0.0) | 45.0 (SE = 8.57; p = 0.174) |
| Water disgust | 0.50 (0.1 – 4.1) | 0.0 (0.0 - 0.35) | 8.0 (SE = 9.04; p = 0.012) |
| Quinine pleasantness | -8.7 (-9.2 – 6.7) | -10 (-10 – -9.5) | 11.5 (SE = 9.19; p = 0.031) |
| Quinine disgust | 8.5 (6.1 – 9.2) | 10 (6.95 – 10) | 44.5 (SE = (9.20; p = 0.174) |
| SDS | 66 (54 – 76) | 61 (54.5 – 69.5) | 27.5 (SE = 9.43; p = 0.681) |
| DSR | 16 (14 – 20) | 15 (10.75 – 20) | 22.0 (SE = 9.42; p = 0.351) |

SDS is Self-Disgust Scale,⁶ DSR is Disgust Scale Revised.^{7,20} In first two columns median with interquartile ranges (in brackets) are given. Last column includes U Mann Whitney results, test standard error (SE) and its associated p value.

TASTE TASK

Participant code: _____
Date: _____

CUP 1

Pleasantness

-10 (very unpleasant)

(very pleasant) +10

Disgust

0 (not disgusting at all) (extremely disgusting) +10

Figure DS1: Visual scales: Response sheet provided to the participants. After taking a sip out of a cup filled with 10 ml. of liquid and maintaining the liquid in their mouths for 10 seconds they had to rate the pleasantness and disgust produced by the intake of the beverage. Pleasantness line measured 20 cm whereas the disgust line measured 10 cm.




Figure DS2 Flow diagram of taste evaluation




Figure DS3 Whole group pleasantness and disgust ratings. The left column shows the histograms of the pleasantness ratings for the 3 conditions in the whole group of participants. The right column shows the histograms of the disgust ratings for the 3 conditions in the whole group of participants.




Figure DS4 Scatter plot with juice pleasantness ratings.




Figure DS5 Scatter plot with quinine pleasantness ratings




Figure DS6 Scatter plot of the disgust ratings for Juice (Y axis) and the Self-Disgust Scale results (X axis) in the Borderline Personality Disorder group.

Additional references

- 13 Steiner JE, Rosenthal-Zifroni A, Edelstein EL. Taste perception in depressive illness. *The Israel annals of psychiatry and related disciplines*. 1969; 7(2): 223-32.
- 14 Amsterdam JD, Settle RG, Doty RL, Abelman E, Winokur A. Taste and smell perception in depression. *Biol Psychiatry*. 1987; 22(12): 1481-5.
- 15 Berlin I, Givry-Steiner L, Lecrubier Y, Puech AJ. Measures of anhedonia and hedonic responses to sucrose in depressive and schizophrenic patients in comparison with healthy subjects. *European psychiatry : the journal of the Association of European Psychiatrists*. 1998; 13(6): 303-9.
- 16 Steiner JE, Lidar-Lifschitz D, Perl E. Taste and odor: reactivity in depressive disorders, a multidisciplinary approach. *Perceptual and motor skills*. 1993; 77(3 Pt 2): 1331-46.
- 17 Potts AJ, Bennett PJ, Kennedy SH, Vaccarino FJ. Depressive symptoms and alterations in sucrose taste perception: cognitive bias or a true change in sensitivity? *Can J Exp Psychol*. 1997; 51(1): 57-60.
- 18 Dichter GS, Smoski MJ, Kampov-Polevoy AB, Gallop R, Garbutt JC. Unipolar depression does not moderate responses to the Sweet Taste Test. *Depress Anxiety*. 2010; 27(9): 859-63.
- 19 Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al. The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry*. 1998; 59 Suppl 20: 22-33;quiz 4-57.
- 20 Olatunji BO, Haidt J, McKay D, David B. Core, animal reminder, and contamination disgust: Three kinds of disgust with distinct personality, behavioral, physiological, and clinical correlates. *J Res Pers*. 2008; 42(5): 1243-59.
- 21 Beck AT, Steer RA, Ball R, Ranieri W. Comparison of Beck Depression Inventories -IA and -II in psychiatric outpatients. *J Pers Assess*. 1996; 67(3): 588-97.
- 22 Hamilton M. Rating depressive patients. *J Clin Psychiatry*. 1980; 41(12 Pt 2): 21-4.
- 23 Dunn OJ. Multiple comparisons using rank sums. *Technometrics*. 1964; 6(3): 241-52.

BJPpsych

The British Journal of Psychiatry

Hedonic and disgust taste perception in borderline personality disorder and depression

Gonzalo Arrondo, Graham K. Murray, Emma Hill, Bence Szalma, Krishna Yathiraj, Chess Denman and Robert B. Dudas

BJP 2015, 207:79-80.

Access the most recent version at DOI: [10.1192/bjp.bp.114.150433](https://doi.org/10.1192/bjp.bp.114.150433)

Supplementary Material

Supplementary material can be found at:
<http://bjp.rcpsych.org/content/suppl/2015/05/14/bjp.bp.114.150433.DC1.html>

References

This article cites 11 articles, 0 of which you can access for free at:
<http://bjp.rcpsych.org/content/207/1/79#BIBL>

Reprints/permissions

To obtain reprints or permission to reproduce material from this paper, please write to permissions@rcpsych.ac.uk

You can respond to this article at

[/letters/submit/bjprcpsych;207/1/79](http://letters.submit/bjprcpsych;207/1/79)

Downloaded from

<http://bjp.rcpsych.org/> on August 10, 2015
Published by The Royal College of Psychiatrists
