Table S3. Overlap of wCNEs with known *cis*-regulatory elements. The genes shown in this table were extracted from literature searches for dissected *cis*-regulatory elements. For each gene, we show the number of spatially associated wCNEs (plus the additional wCNE associated with *lir-1*), the number of wCNEs that overlap dissected *cis*-elements and the references in which the experiments are described. In the case of *lin-26*, there is one wCNE, which is closer to the *lir-1* transcription start site than *lin-26*, so it was associated with *lir-1* in our dataset, however experiments showed that it is likely to be a *lin-26* distal *cis*-element.

Gene	Number of wCNEs	Number of wCNEs that overlap experimentally tested sequences showing cis-regulatory activity	Ref.
myo-2	1	1	[1]
lin-26	3 (+ 1 spatially associated with lir-1)	3 (+1)	[2]
ceh-22	2	1 (the other wCNE is intronic)	[3]
ser-2	5	5	[4]
ric-4	1	1	[5]
egl-5	1	0 (wCNE lies downstream)	[6]
ttx-3	0	-	[4]
sra-11	0	-	[4]
kal-1	0	-	[4]
hen-1	0	-	[4]
unc-17	0	-	[4]
ceh-10	0	-	[4]
ceh-23	0	-	[4]
dpy-7	0	-	[7]
egl-17	0	-	[8]
mec-3	0	-	[9]
her-1	0	-	[10]

References

- Kalb JM, Beaster-Jones L, Fernandez AP, Okkema PG, Goszczynski B, McGhee JD: Interference between the PHA-4 and PEB-1 transcription factors in formation of the Caenorhabditis elegans pharynx. J Mol Biol 2002, 320(4):697-704.
- 2. Landmann F, Quintin S, Labouesse M: Multiple regulatory elements with spatially and temporally distinct activities control the expression of the epithelial differentiation gene lin-26 in C. elegans. *Dev Biol* 2004, 265(2):478-490.
- 3. Vilimas T, Abraham A, Okkema PG: An early pharyngeal muscle enhancer from the Caenorhabditis elegans ceh-22 gene is targeted by the Forkhead factor PHA-4. *Dev Biol* 2004, **266**(2):388-398.
- 4. Wenick AS, Hobert O: Genomic cis-regulatory architecture and trans-acting regulators of a single interneuron-specific gene battery in C. elegans. *Dev Cell* 2004, **6**(6):757-770.
- 5. Hwang SB, Lee J: Neuron cell type-specific SNAP-25 expression driven by multiple regulatory elements in the nematode Caenorhabditis elegans. *J Mol Biol* 2003, **333**(2):237-247.
- 6. Teng Y, Girard L, Ferreira HB, Sternberg PW, Emmons SW: **Dissection of cisregulatory elements in the C. elegans Hox gene egl-5 promoter**. *Dev Biol* 2004, **276**(2):476-492.
- 7. Gilleard JS, Barry JD, Johnstone IL: cis regulatory requirements for hypodermal cell-specific expression of the Caenorhabditis elegans cuticle collagen gene dpy-7. *Mol Cell Biol* 1997, **17**(4):2301-2311.
- 8. Cui M, Han M: Cis regulatory requirements for vulval cell-specific expression of the Caenorhabditis elegans fibroblast growth factor gene egl-17. *Dev Biol* 2003, 257(1):104-116.
- 9. Xue D, Finney M, Ruvkun G, Chalfie M: **Regulation of the mec-3 gene by the C.elegans homeoproteins UNC-86 and MEC-3**. *Embo J* 1992, **11**(13):4969-4979.
- 10. Li W, Streit A, Robertson B, Wood WB: Evidence for multiple promoter elements orchestrating male-specific regulation of the her-1 gene in Caenorhabditis elegans. *Genetics* 1999, **152**(1):237-248.