Dalton Transactions

ARTICLE

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2014, Accepted 00th January 20XX

DOI: 10.1039/x0xx00000x

www.rsc.org/

Multiple Deprotonation of Primary Aromatic Diamines by LiAlH₄

Robert J. Less,^{*,a} Lucy. K. Allen,^a Alexander Steiner^b and Dominic S. Wright^a

Reaction of LiAlH₄ with 1,2-phenylenediamine (1H₄) in THF results in formation of the metallocyclic amido-/imido complex [{Al(1H₂)}₂{Al(1H₂)}₂[Li(THF)₂]₄ (3), while in the presence of various Lewis base ligands 1,8-diaminonaphthalene (2H₄) gives the amido-('ate') complexes [Al(2H₂)₂]⁻[Li(LL')]⁺ [L = THF, L' = PMDETA (*N*,*N*,*N'*,*N''*-pentamethyldiethylenetriamine) (4); L = L' = TMEDA (*N*,*N*,*N'*,*N'*-Tetramethylethylenediamine) (5)]. The latter complexes provide evidence of intermediates in the proposed reaction pathway for formation of the cyclic framework of the tetraanion [{Al(1H₂)₂{Al(1H)₂}₂]⁴⁻ of 3.

1. Introduction

Group 13 amido (R_2N^{-}) and imido (RN^{2-}) compounds are important classes of main group metal nitrogen compounds owing to their potential use in a number of synthetic and technologically-important applications, such as reagents for organic and inorganic transformations and single-source precursors for technologically important III/V semiconductor materials (*e.g.*, AlN and GaN).¹ More recent developments in lithium amido aluminate chemistry include derivatives based on TMP (2,2,6,6-tetramethylpiperidide) as reagents to effect the metalation of aromatic compounds.² Group 13 *amido* (R_2N^{-}) and *imido* (RN^{2-}) compounds have been obtained using a broad range of synthetic routes.³

One interesting class of these compounds are alkali metal primary-amido aluminium 'ate' complexes containing $[Al(NHR)_4]^-$ anions,⁴ which can be employed as precursors for imido complexes via deprotonation of the Al-bonded RNHgroups with strong organometallic bases such as "BuLi.⁵ This approach to imido aluminium compounds can, however, suffer from the drawback that the NH protons may not be reactive enough to effect deprotonation, with the result that the RNH functionality of the precursor is retained and the "BuLi base molecules are simply incorporated into the resulting cage structure.⁶ Alternatively, separate studies by Roesky, Chivers and Nöth have shown that direct reactions of lithium and sodium aluminium hydride (MAlH₄; M = Li or Na) with secondary (R₂NH) and mono-functional primary amines (RNH₂) can be employed to produce a range of amido- and mixed amido/imido lithium 'ate' compounds.7

Our current focus on this area stems jointly from long standing interests in the synthesis and structural chemistry of main group metal imido complexes and their phosphanediide (RP²⁻) counterparts⁸ and from a more recent realization that key reagents which we have employed in their synthesis can be used to effect stoichiometric and catalytic dehydrocoupling of a range of bonds.⁹ Pertinent to the current study, we have found that the reaction of the redox-active superbase mixture Sb(NMe₂)₃/ⁿBuLi with the 1,2-(PH₂)₂C₆H₄ results in the formation of the benzo-1,2,3-triphosphilide anion $[C_6H_4P_3]$ via a process involving a stepwise oxidative coupling of the P-H bonds (Scheme 1a), in combination with the reduction of Sb^{III} to a Zintl compound.¹⁰ Remarkably, the same type of coupling occurs in the case of the reaction of the diamine analogue ophenylene diamine, $1,2-(NH_2)_2C_6H_4$, with the superbase mixture Sn(NMe₂)₂/ⁿBuLi at elevated temperature, which gives the benzo-1,2,3-triazolyl anion and Sn metal (Scheme 1b).¹¹





Scheme 1 Reactivity of $1,2\text{-}C_6\text{H}_4(\text{EH}_2)_2$ (E = N, P) with p-block metal bases and "BuLi.

Such N-H/N-H coupling is extremely rare even in the case of transition metal-mediated reactions and may have broader scope in the synthesis of a range of nitrogen heterocycles.¹²

Most recently it was found that amides, like $Al(NMe_2)_3$, and hydrides, such as ('BuO) AlH_2 and Li AlH_4 , are effective precatalysts for N-H/B-H dehydrocoupling.¹³ With this background in mind, we focus in the current study on the reactions of the most readily available reagent of this class (Li AlH_4) with aromatic diamines, with a view to exploring whether this simple reagent might also be capable of N-H/N-H dehydrocoupling.

2. Results and Discussion

Following on from previous studies in this area, we selected 1,2-phenylenediamine (1H₄) and 1,8-diaminonaphthalene (2H₄) as substrates for the current investigations (Fig. 1). Although 1H₄ and 2H₄ and anions derived from their deprotonation have been used extensively as ligands for a large number of transition metal complexes,^{14,15} only three examples of Group 13 complexes containing the doubly-deprotonated $[1H_2]^{2-}$ ligand are known (*i.e.*, deprotonated at each of the NH₂ groups, see Scheme 2).¹⁶ In addition, only one example of a main group metal complex containing the $[2H_2]^{2-}$ has been structurally characterized previously (*i.e.*, again with deprotonation at each of the NH₂ groups, see Scheme 3).¹⁷



Fig. 1 Structures of the precursor diamines used in the current work $1H_4$ and $2H_4$.

Addition of solid LiAlH₄ to solutions of the diamines $1H_4$ or 2H₄ using various stoichiometries in THF results in rapid exothermic reactions with the visible evolution of H₂ gas and the formation of colourless to light yellow solutions. Particular care had to be taken to ensure the rigorous exclusion of air from these reactions as the solutions generated react instantaneously with traces of atmospheric oxygen to give blue, purple or brown solutions (depending on the degree of contamination). Initially, a 1:1 ratio of the diamines to LiAlH₄ was employed in trial reactions, with the intention of fully deprotonating both amine groups to give the tetraanions $[1]^{4-}$ and $[2]^{4-}$. Unfortunately, in both cases the products of these reactions could not be isolated despite numerous attempts at crystallization. However, using a reduced ratio of LiAlH₄ with $1H_4$ (2 : 3, respectively) in THF allowed the isolation of highly air-sensitive colourless crystals of the new aluminium complex $[{Al(1H_2)}_2{Al(1H)_2}_2][Li(THF)_2]_4$ (3) in 48% yield, containing doubly and triply-deprotonated $1{H_2}^{2-}$ and $1{H}^{3-}$ anions (Scheme 2).

The room-temperature ¹H NMR spectrum of crystalline **3** in D₈-THF (ESI) shows two distinct sets of aromatic C-H resonances for the $[1H]^{3-}$ [δ 6.16 (s) and 6.08 (s)] and $[1H_2]^{2-}$ [δ 5.0-6.5 (br. mult.)] ligands in the correct 1:1 ratio. However, the

NH protons of these anions are indistinguishable and only appear as a broad singlet (δ 3.08 ppm), but with the correct 24:8 ratio with respect to the total aromatic C-H protons. Reducing the temperature failed to resolve these resonances, possibly implying fast scrambling of the H-atoms between the [1H]³⁻ and [1H₂]²⁻ ligands.

$$1 \text{ LiAIH}_4 + 6 [1H_4] \longrightarrow [{Al(1H_2)}_2{Al(1H)_2}_2][Li(THF)_2]_4 + 16 \text{ H}_2$$



Scheme 2 The stoichiometry of the reaction producing 3, showing the structure of the complex and the observed deprotonation pattern of the $[1\rm H_4]$ substrate.

A further feature of the ¹H NMR spectra of samples of **3**, which were dried under vacuum prior to transfer to a glove box for analysis, is the lower than expected integral of the THF ligands. This feature is probably largely responsible for the higher than expected N analysis found for isolated samples of the complex (found 14.3% *vs.* 12.5 %; although the C and H analyses were acceptable).

Double- and single-deprotonation of the NH₂ groups of monofunctional aliphatic and aromatic primary amines has been seen before using LiAlH₄.^{7b,c} An example is the reaction of DippNH₂ (Dipp = $2,6^{-i}$ PrC₆H₃) which gives the dimer [Al{(DippNH)₂}₂(μ -NDipp)][Li(THF)]₂.^{7b} However, to the best of our knowledge the current study represents the first investigation of the reactions of LiAlH₄ with aromatic diamines. The formation of the triply-deprotonated [1H]³⁻ trianion in **3** is similar to that seen in the reaction of 1H₄ with Sn(NMe₂)₂/^{*n*}BuLi at room temperature,¹¹ while the quadruple-deprotonation of 1H₄ (to [1]⁴⁻) only appears to occur in the presence of the more basic reagent mixture Sb(NMe₂)₃/^{*n*}BuLi.¹⁸

Reactions of 2H₄ with LiAlH₄ proceeded in a similar manner to those with 1H4, but the products appeared to be much less sensitive to traces of oxygen. Although a variety of reaction conditions and different reactant stoichiometries were employed, only products resulting from double deprotonation of 2H₄ could be isolated and only in the presence of chelating donor ligands. The new complexes $[Al(2H_2)_2]^{-1}$ $[Li(PMDETA)(THF)]^+$ (4) and $[Al(2H_2)_2]^-[Li(TMEDA)_2]^+$ (5) are obtained in highest yields (48% and 53%, respectively) from the 1:2 stoichiometric reactions of LiAlH₄ with 2H₄ in THF, in presence of an excess of the Lewis base donor ligands PMDETA and TMEDA (respectively) (Scheme 3). Unlike the room temperature ¹H NMR spectrum of **3**, the ¹H NMR spectra of **4** and **5** are well resolved in the aromatic region of the $[2H_2]^{2-}$ dianion ligands (ESI). The 1,8-naphthyl groups in both appear in the range δ 5.8-6.6, with the N-H protons being found at δ 3.8. The ¹H NMR spectra of **5** also show that a significant amount of residual THF is preserved in crystalline samples of the complex after isolation under vacuum. Both C, H and N analysis and integration of the ¹H NMR spectrum show that there are *ca.* three TMEDA molecules per THF normally found in solid samples (*i.e.*, 1 molecule of THF for every 1.5 molecules of **5**).

 $LiAIH_4 + 2[\mathbf{2}H_4] \xrightarrow{L/L'} [AI(\mathbf{2}H_2)_2]^{-}[Li(L)(L')]^{+} + 4H_2$ $L = PMDETA; L' = THF (\mathbf{4})$ $L = L' = TMEDA (\mathbf{5})$



Scheme 3 The reactions producing the related ion-separated complexes 4 and 5, showing the structure of the aluminate anion and the deprotonation pattern of $2H_4$.

1,8-diaminonaphthalene ($2H_4$), particularly in its doubly deprotonated form $[2H_2]^{2-}$, has been used extensively as a ligand in a number of transition metal complexes.¹⁵ However, although a large number of complexes containing *N*,*N*-disubstituted ligands of the type $[2R_2]^{2-}$ (R = alkyl, aryl or silyl) have been reported for main group metals,¹⁹ main group metal complexes containing the parent $[2H_2]^{2-}$ ligand are very rare¹⁷ and there are no previously reported examples of Group 13 metal complexes containing the $[2H_2]^{2-}$ ligand.

Single-crystal X-ray diffraction studies of the new complexes **3**, **4** and **5** were undertaken at 180(2) K. Details of the structure refinements and data collections on **3** and **4** are collected in Table 1 (Experimental Section). Since the X-ray data for **5** was of poor quality [with lowest obtained R(1) = 10.34%, despite repeated data collections] the structure of the complex is not discussed in detail here (see ESI).

 $[{Al(1H_2)}_2{Al(1H)_2}_2][Li(THF)_2]_4$ (3) crystallizes in monoclinic space group $P2_1/n$ and its molecular structure is illustrated in Fig. 2. The complex has an ion-paired structure resulting from the association of an aluminate $[{Al(1H_2)}_2{Al(1H)}_2]^{4-}$ tetraanion with four $[Li(THF)_2]^+$ cations. A schematic of the structure is shown in the insert to Scheme 2. The $[{Al(1H_2)}_2{Al(1H)}_2]^{4-}$ tetraanion is constructed (formally) from two $[Al(1H)_2]^{3-}$ and $[Al(1H_2)]^+$ subunits which form a cyclic arrangement composed of a central eight-membered (*twist-chair*²⁰) Al_4N_4 ring unit, using the imido-N atoms of the $[Al(1H)_2]^{3-}$ fragments.



Fig. 2 (a) Molecular structure of $[{Al(1H_2)}_2{Al(1H)}_2]_2[Li(THF)_2]_4$ (**3**). C-bonded Hatoms and the disordered components of the THF ligands have been omitted for clarity. Selected bond lengths (Å) and angles (°): Al(1)-N(1) 1.846(3), Al(1)-N(2) 1.858(4), Al(1)-N(3) 1.843(3), Al(1)-N(4) 1.865(3), Al(3)-N(7) 1.869(3), Al(3)-N(8) 1.831(5), Al(3)-N(9) 1.880(3), Al(3)-N(10) 1.832(5), Al(2)-N(3) 1.818(3), Al(2)-N(5) 1.880(3), Al(2)-N(6) 1.874(3), Al(2)-N(7) 1.845(3), Al(4)-N(1) 1.826(3), Al(4)-N(9) 1.862(3), Al(4)-N(11) 1.880(3), Al(4)-N(12) 1.877(3), Li-N range 2.15(1)-2.25(1), N-Al-N (Al_4N_4 ring) range 117.3(2)-124.9(2), N-Al-N (chelate) range 87.2(9)-90.0(2), Al-N-Al (Al_4N_4 ring) range 119.4(2)-122.0(2); (b) The core structure of the complex, with the THF ligands and aromatic rings removed.



This arrangement results in two distinct Al-environments [Al(1)/Al(3) and Al(2)/Al(4)], each of which have pseudo-tetrahedral metal geometries with the Al centres being bonded

solely to N atoms. The Al-N bond lengths within **3** are in the range 1.818(3)-1.880(3) Å, with no significant differences between the amido N(-H)-Al and imido N-Al bonds being discernible.²¹ The tetraanion of **3**, containing four Al-atoms, is the largest species of this type to be reported. Previously, amido/imido-aluminate anions containing a maximum of two Al atoms have been observed.^{7b,c}

A surprising feature of the structural arrangement of **3** is the presence of two Li⁺ coordination environments. Although all of the Li⁺ cations are pseudo-tetrahedral, Li(1) and Li(4) are each bound to one amido N(H) atom of a $[Al(1H)_2]^{3-}$ and one of a $[Al(1H_2)]^+$ fragment, while Li(2) and Li(3) are bonded to the imido-N atom of a $[Al(1H)_2]^{3-}$ unit and an amido N(H) atom of a $[Al(1H_2)]^+$ unit. The N-Li bond lengths are in the range 2.15(1) - 2.25(1) Å, within the normal range observed previously.²¹ The reason for the unsymmetrical coordination of the Li⁺ cations in **3** can be traced to the disposition of the C₆H₄ rings of the $[Al(1H)_2]^{3-}$ subunits at either end of the $[{Al(1H_2)}_2{Al(1H)}_2]_2^{4-}$ tetraanion. As can be seen from Fig. 2b, showing the core structure of the complex, the $[Al(1H_2)]^{3-}$ subunits have virtual mirror symmetry with respect to the central Al₄N₄ ring, which limits the availability of the imido-N atoms to coordinate all of the Li⁺ cations.



Fig. 4 Structure of (a) the anions and (b) the cation of $[Al(2H_2)_2]^{-1}$ [Li(PMDETA)(THF)]^{*} (**4**). C-bonded H-atoms have been omitted for clarity. Bound H atoms have been omitted for clarity. Selected bond lengths (Å) and angles (°): Al(1)-N(1) 1.821(3), Al(1)-N(2) 1.814(3), Al(1)-N(3) 1.831(3), Al(1)-N(4) 1.815(3), N-Al-N (chelate) range 95.2(1)-95.6(1), N-Al-N (between $[2H_2]^{2^{-1}}$ ligands) range 116.2(2)-118.3(2), C-N-Al range 127.6(2)-129.6(3).

 $[Al(2H_2)_2]^{-}[Li(PMDETA)(THF)]^{+}$ (4) crystallizes in monoclinic space group C2/c and its molecular structure is illustrated in Fig. 4. The complex is ion-separated and consists of $[Li(PMDETA)(THF)]^+$ cations and $[Al(2H_2)_2]^-$ anions. The structure of $[Al(2H_2)_2]^{-}[Li(TMEDA)_2]^{+}$ (5) (see ESI) only differs in the nature of the cation present. The formation of the $[Al(2H_2)_2]^-$ anion in 4 and 5 is similar to the outcome of the reactions of primary amines (RNH_2) with $MAlH_4$ (M = Li, Na)(1:4 equivalents) which give $[Al(NHR)_4]^-$ ions.^{7c} The Al³⁺ cation of the anion of 4 has a highly distorted tetrahedral geometry, resulting from the sharp angles within the chelate rings [N-Al-N range 95.2(1) - 95.6(1)°; cf. range 116.2(2) -118.3(2)° for the remaining N-Al-N angles]. The sharp chelate angles and the Al-N bond lengths [range 1.814(3) - 1.831(3) Å] are similar to those found in 3. Despite the adoption of planar sp^2 hybridization at the N-atoms, there is no evidence for significant double-bond character in the Carvel-N bonds, whose bond lengths are typical of those found in aromatic amines [range 1.359(5) - 1.384(5) Å]. Its simplicity apart, to our knowledge the anion of 4 is the only example of a mononuclear metal complex containing the $[2H_2]^{2-}$ ligand. In the context of the current study the $[Al(2H_2)_2]^{-1}$ anion can be seen as representing a model precursor for the [Al(1H)₂]³⁻ units found in complex 3, resulting from further double deprotonation (see the conclusions section of this paper).

3. Conclusions

We have presented here the first study of the reactions of LiAlH₄ with primary diamines. The new compound $[{Al(1H_2)}_2{Al(1H)}_2][Li(THF)_2]_4$ (3) is especially interesting as contains the largest of this class of amido/imido aluminate anions, the tetraanion $[{Al(1H_2)}_2{Al(1H)}_2]^{4^-}$.



Scheme 4 Pror	osed reaction	nathway	of formation	of the tetraar	nion of 3
Scheme 4 Frop	Juseu reaction	patriway	or rormation		11011 UI 3 .

The formation of the mononuclear anions in $[Al(2H_2)_2]^{-1}$ $[Li(PMDETA)(THF)]^+$ (4) and $[Al(2H_2)_2]^-[Li(TMEDA)_2]^+$ (5) follows a more straightforward pathway that is similar to that seen previously in certain reactions of MAlH₄ (M = Li, Na).⁷ The $[Al(2H_2)_2]^-$ anions in the latter give a clue to the way in which the tetraanion of 3 is formed (Scheme 4). It is likely, given the observed (2:3) stoichiometry of the reaction, that the AlH_4 ion will react to give a mixture of $[H_2Al(1H_2)]$ and $[Al(1H_2)_2]$ in the initial stage, which then further to give the anion of 3 directly. Finally, we have found in the current study that 3, 4 and 5 are thermally stable at reflux in THF and do not appear to undergo oxidative coupling in a similar manner to that found previously in the case of $1H_4$ with Sn(NMe₂)₂/ⁿBuLi.¹¹ The reason for this is almost certainly the lack of a redox active metal species which will affect the N-H/N-H dehydrocoupling step.

Acknowledgements

We thank the EU (ERC Advanced Investigator Grant for D.S.W.), the EU (R.J.L. and L.K.A.). We also thank Dr. J. E. Davies for collecting X-ray data on **3**, **4** and **5**.

Notes and references

^a Department of Chemistry, Lensfield Road, Cambridge CB2 1EW, United Kingdom.

^b University of Liverpool, Department of Chemistry, Crown Street, Liverpool L69 7ZD, United Kingdom.

Electronic Supplementary Information (ESI) available: crystal and NMR data for compounds **3-5**. See DOI: 10.1039/b000000x/

Experimental

General Experimental Procedures

All compounds described in this paper are air and moisture sensitive. Preparations were performed on a double-manifold vacuum line under a nitrogen atmosphere. The products were isolated and stored with the aid of a nitrogen-filled glove box (Saffron type b), equipped with Cu and molecular sieve columns in order to remove O₂ and moisture, respectively. TMEDA and PMDETA (Aldrich) were distilled from Na and stored under argon. All ¹H and ¹³C NMR spectra were recorded using a Brüker DPX 500 MHz NMR spectrometer. Elemental (C, H, N) analyses were obtained using an Exeter CE-440 Elemental Analyser. Samples for analysis (1-2 mg) were placed in pre-weighed, airtight aluminium boats in the glove box prior to analysis.

Synthesis of $[{Al(1H_2)}_2 {Al(1H)}_2]_2 [Li(THF)_2]_4$ (3): LiAlH₄ (23) mg, 0.61 mmol) was added to a solution of 1,2phenylenediamine (100 mg, 0.92 mmol) in 5 ml THF and the mixture stirred for 16h at room temperature. The solution was filtered into a crystallization tube (20 mm diameter) and the volume of solvent reduced in vacuo to approximately 1 ml. Hexane (10 ml) was then carefully layered onto the solution and the tube was sealed and allowed to remain undisturbed for one week after which time the product had solidified. The solvents were then decanted off and the solids collected and allowed to dry in the glove box. Yield 99 mg (48 %). Found (%): C 61.3, H 7.0, N 14.3; Calcd. for C₆₈H₉₆Al₄Li₄N₁₂O₈: C 60.7, H 7.2, N 12.5. ¹H NMR (400.1 MHz, 25 °C, D₈-THF), $\delta/\text{ppm} = 6.13$ (br s, 8H, $[ArH(NH)(N)]^{3-}$), 6.05 (br s, 8H, $[ArH(NH)(N)]^{3-}$, 5.9-6.5 (br m, 8H, $ArH(NH)_2^{2-}$), 3.60 (m, THF), 3.05 (br s, 8H, NH), 1.77 (m, THF); ¹³C{¹H} NMR (400.1 MHz, 25 °C, D8-THF), δ/ppm = 145.5 (br m, ArH), 114.5 (br m, ArH), 110.8 (br m, ArH), 68.2 (m, THF), 26.4 (THF).

Synthesis of $[Al(2H_2)_2]^{-}[Li(PMDETA)(THF)]^{+}$ (4): LiAlH₄ (12) mg, 0.32 mmol) was added to a solution of 1,8diaminonaphthalene (100 mg, 0.63 mmol) in 5 ml THF containing 0.5 ml PMDETA (2.4 mmol). The mixture was heated to reflux for 1h and sufficient hot THF (approx. 3 ml) was added to dissolve the precipitate which had formed. The solution was filtered whilst hot and allowed to cool slowly to room temperature resulting in crystallization of the product. The mother liquor was decanted off and the crystals allowed to dry in the glove box. Yield 90 mg (48 %). Found (%): C 67.0, H 7.9, N 16.6; Calcd. for C₃₃H₄₇AlLiN₇O: C 67.0, H 8.0, N 16.6. ¹H NMR (400.1 MHz, 25 °C, D_8 -THF), δ /ppm = 6.62 (t, *J* = 8.0 Hz, 4H, ArH), 6.33 (dd, *J* = 8.0, 1.2 Hz, 4H, ArH), 5.86 (dd, J = 8.0, 1.2 Hz, 4H, ArH), 3.81 (s, 4H, ArNH), 3.61 (m,4H, THF), 2.39 (td, J = 6.8, 0.8 Hz, 4H, PMDETA CH₂), 2.28 (td, J = 6.8, 0.8 Hz, PMDETA CH₂), 2.19 (s, 3H, PMDETA CH₃), 2.14 (s, 12H, PMDETA CH₃). 1.76 (m, 4H, THF); ¹³C{¹H} NMR (400.1 MHz, 25 °C, D8-THF), δ /ppm = 155.8

(Ar), 140.0 (Ar), 126.8 (Ar), 118.5 (Ar), 112.1 (Ar), 107.3 (Ar), 68.2 (THF), 58.7 (PMDETA), 56.8 (PMDETA), 46.1 (PMDETA), 43.5 (PMDETA), 26.4 (THF).

Synthesis of $[Al(2H_2)_2]^{-}[Li(TMEDA)_{1,5}(THF)_{0,5}]^{+}$ (5): LiAlH₄ (12 mg, 0.32 mmol) was added to a solution of 1,8diaminonaphthalene (100 mg, 0.63 mmol) in 5 ml THF containing 0.5 ml TMEDA (3.6 mmol). The mixture was heated to reflux for 1h, filtered and the volume of solvent reduced in vacuo whilst hot until the onset of crystallization occurred. The solution was allowed to cool to room temperature to complete the crystallization whereupon the mother liquor was decanted off and the product collected and allowed to dry in the glove box. Yield 100 mg (53 %). Found (%): C 66.7, H 8.2, N 17.6; Calcd. for C₃₂H₄₈AlLiN₈·C₃₀H₄₀AlLiN₆O C 66.9, H 8.0, N 17.6. ¹H NMR (400.1 MHz, 25 °C, D8-THF), δ /ppm = 6.63 (t, *J* = 7.6 Hz, 4H, ArH), 6.34 (dd, *J* = 7.6, 0.8 Hz, 4H, ArH), 5.86 (dd, J = 7.6, 0.8 Hz, 4H, ArH), 3.82 (s, 4H, ArNH), 3.61 (m, THF), 2.30 (8H, s, TMEDA CH₂), 2.15 (s, 24H, TMEDA CH₃), 1.77 (m, THF); ¹³C{¹H} NMR (400.1 MHz, 25 °C, D8-THF), δ/ppm = 155.8 (Ar), 140.0 (Ar), 126.8 (Ar), 118.5 (Ar), 112.1 (Ar), 107.3 (Ar), 68.2 (THF), 58.9 (TMEDA), 46.2 (TMEDA), 26.4 (THF).

X-Ray Crystallography

Data for all complexes were collected on a Nonius KappaCCD diffractometer and solved by direct methods and refined by full-matrix least squares on F^2 . All of the THF molecules in the structure of **3** [except those attached to Li(4)] exhibit site disorder in which the C-atoms of the backbone of the ligands were modelled over two 50:50 sites.

Compound	3	4	
Chemical formula	$\begin{array}{c} C_{68}H_{96}Al_4Li_4N_{12}O_8 \\ 0.5(C_4H_8O) \cdot 0.5(C_6H_{14}) \end{array}$	$\frac{C_{20}H_{16}AlN}{C_{13}H_{31}LiN_3O} +$	
FW / g⋅mol ⁻¹	1424.39	591.70	
Crystal system	Monoclinic	Monoclinic	
Space group	$P2_1/n$	C2/c	
<i>a /</i> Å	15.8568(5)	34.6994(4)	
<i>b</i> / Å	25.4155(9)	14.6711(2)	
<i>c /</i> Å	20.1493(8)	30.4757(4)	
β/°	101.022(2)	120.945(1)	
$V / Å^3$	7970.5(5)	13306.2(3)	
Ζ	4	16	
$ ho_{ m calcd}/ m Mg{\cdot}m^{-3}$	1.187	1.181	
μ (MoK α) / mm ⁻¹	0.118	0.097	
Reflections collected	11879	12123	
Independent reflections	7865	6898	
$R_{\rm int}$	0.042	0.097	
$R_1, WR_2 [I > 2\sigma(I)]$	0.067, 0.157	0.081, 0.238	
R_1 , w R_2 (all data)	0.108, 0.181	0.127, 0.261	

Data in common $\lambda = 0.71070$ nm, T = 180(2)K.

- D. A. Neumayer and J. G. Ekerdt, *Chem. Mater.*, 1996, 8, 9; A. C. Jones, *Chem. Soc. Rev.*, 1997, 101.
- H. Naka, J. V. Morey, J. Haywood, D. J. Eisler, M. McPartlin, F. García, H. Kudo, Y. Kondo, M. Uchiyama and A. E. H. Wheatley, *J. Am. Chem. Soc.* 2008, **130**, 16193; D. R. Armstrong, E. Crosbie, E. Hevia, R. E. Mulvey, D. L. Ramsay and S. D. Robertson, *Chem. Sci.*, 2014, **5**, 3031.
- M. Veith, *Chem. Rev.*, 1990, **90**, 3; S. J. Schauer and G. H. Robinson, *J. Coord. Chem.*, 1993, **30**, 197; S. Schulz, L. Haming, R. Herbst-Irmer, H. W. Roesky and G. M. Sheldrick, *Angew. Chem. Int. Ed. Engl.*, 1994, **33**, 969; J. K. Brask and T. Chivers, *Angew. Chem. Int. Ed.*, 2001, **40**, 3960; G. M. Aspinall, M. C. Copsey, A. P. Leedham and C. A. Russell, *Coord. Chem. Rev.*, 2002, **227**, 217.
- 4 M. A. Beswick, N. Choi, C. H. Harmer, M. McPartlin, M. E. G. Mosquera, P. R. Raithby, M. Tombul and D. S. Wright, *Chem. Commun.*, 1998, 1383; J. S. Silverman, C. J. Carmalt, D. A. Neumayer, A. H. Cowley, B. G. McBurnett and A. Decken, *Polyhedron*, 1998, **17**, 977.
- 5 D. A. Atwood and D. Rutherford, *Organometallics*, 1996, **15**, 436; M. C. Copsey, J. C. Jeffery, C. A. Russell, J. M. Slattery and J. A. Straughan, *Chem. Commun.*, 2003, 2356; D. Rutherford and D. Atwood, *J. Am. Chem. Soc.*, 1996, **118**, 11535.
- 6 J. K. Brask, T. Chivers and G. P. A. Yap, *Chem. Commun.*, 1998, 2543; J. K. Brask, T. Chivers, G. Schatte and G. P. A. Yap, *Organometallics*, 2000, **19**, 5683.
- 7 (a) G. Linti, H. Nöth and P. Rahm, Z. Naturforsch, 1988, 43b, 1101; (b) M. L. Monterro, H. Wessel, H. W. Roesky, M. Teichert and I. Usón, Angew. Chem. Int. Ed. Engl., 1997, 36, 629; (c) D. A Eisler and T. Chivers, Can. J. Chem., 2006, 84, 443.
- 8 M. A. Beswick and D. S. Wright, Coord. Chem. Rev., 1998, 176, 373.
- 9 R. J. Less, R. Melen, V. Naseri and D. S. Wright, *Chem. Commun.*, 2009, 4929; R. J. Less, R. L. Melen and D. S. Wright, *RSC Advances*, 2012, **2**, 2191; V. Naseri, R. J. Less, M. McPartlin, R. E. Mulvey and D. S. Wright, *Chem. Commun.*, 2010, **46**, 5000.
- F. García, R. J. Less, V. Naseri, M. McPartlin, J. M. Rawson, M. S. Tomas and D. S. Wright, *Chem. Commun.*, 2008, 859; R. Edge, R. J. Less, V. Naseri, E. J. L. McInnes, R. E. Mulvey and D. S. Wright, *Dalton Trans.*, 2008, 6454.
- 11 R. J. Less, V. Naseri, M. McPartlin and D. S. Wright, *Chem. Commun.*, 2011, **47**, 6129.
- A. Grirrane, A. Corma and H. García, *Science*, 2008, **322**, 1661; C. Zhang and N. Jiao, *Angew. Chem. Int. Ed.*, 2010, **49**, 6174.
- H. J. Cowley, M. S. Holt, R. L. Melen, J. M. Rawson and D. S. Wright, *Chem. Commun.*, 2011, **47**, 2682; M. M. Hansmann, R. L. Melen and D. S. Wright, *Chem. Sci.*, 2011, **2**, 1554; R. J. Less, H. R. Simmonds, S. B. J. Dane and D. S. Wright, *Dalton Trans.*, 2013, **42**, 6337; R. J. Less, H. R. Simmonds and D. S. Wright, *Dalton Trans.*, 2014, **43**, 5785.
- A. Mederos, S. Domínguez, R. Hernández-Molina, J. Sanchiz and F. Brito, *Coord. Chem. Rev.*, 1999, **193-195**, 857; A. Mederos, S. Domínguez, R. Hernández-Molina, J. Sanchiz and F. Brito, *Coord. Chem. Rev.*, 1999, **193-195**, 913.
- 15 E. Sola, F. Torres, M. V. Jiménez, J. A. López, S. E. Ruiz, F. J. Lahoz, A. Elduque and L. A. Oro, J. Am. Chem. Soc., 2001, **123**, 11925; M. V.

Jiménez, E. Sola, J. A. López, F. J. Lahoz and L. A. Oro, Chem. Eur. J., 1998, 4, 1398; M. V. Jiménez, E. Sola, J. Caballero, F. J. Lahoz and L. A. Oro, Angew. Chem. Int. Ed., 2002, 41, 1208; Z. Chen, M. Zeng, Y. Zhang, Z. Zhang and F. Liang, Appl. Organometal. Chem., 2010, 24, 625; J. A. Cabeza, H. Nöth, M. de J. Rosales-Hoz and G. Sánchez-Cabrera, Eur. J. Inorg. Chem., 2000, 2327; L. A. Oro, M. J. Fernandez, J. Modrego, C. Foces-Foces and F. H. Cano, Angew. Chem. Int. Ed. Engl., 1984, 23, 913; H. Matsuzaka, T. Kamura, K. Ariga, Y. Watanabe, T. Okubo, T. Ishii, M. Yamashita, M. Kondo and S. Kitagawa, Organometallics, 2000, 19, 216; M. V. Jiménez, E. Sola, M. A. Egea, A. Huet, A. C. Francisco, F. J. Lahoz and L. A. Oro, Inorg. Chem., 2000, 39, 4868; J. A. Cabeza, J. M. Fernández-Colinas, S. García-Granda, V. Riera and J. F. Van der Maelen, Inorg. Chem., 1992, 31, 1233; C. Nachtigal, S. Al-Gharabli, K. Eichele, E. Lindner and H. A. Mayer, Organometallics, 2002, 21, 105; J. A. Cabeza, J. M. Fernandez-Colinas, V. Riera, M. A. Pellinghelli and A. Tiripicchio, J. Chem. Soc. Dalton Trans., 1991, 371.

- 16 H. Rahbarnoohi, R. L. Wells and A. L. Rheingold, *Chem. Commun.*, 1996, 2661; R. L. Wells, H. Rahbarnoohi, P. B. Glaser, L. M. Liable-Sands and A. L. Rheingold, *Organometallics*, 1996, **15**, 3204; X. Ma, P. Hao, J. Li, H. W. Roesky and Z. Yang, *Z. Anorg. Allg. Chem.*, 2013, **639**, 493.
- 17 W. Clegg, L. Horsburgh, R. E. Mulvey and R. Rowlings, *Chem. Commun.*, 1996, 1739.
- 18 F. García, R. J. Less, V. Naseri, M. McPartlin, J. M. Rawson and D. S. Wright, *Dalton Trans.*, 2008, 997.
- 19 For examples containing aluminium, see T. Riehm, H. Wadepohl and L. H. Gade, *Inorg. Chem.*, 2008, 47, 11467; N. Lavoie, T.-G. Ong, S. I. Gorelsky, I. Korobkov, G. P. A. Yap and D. S. Richeson, *Organometallics*, 2007, 26, 6586; Z. Yang, X. Ma, H. W. Roesky, Y. Yang, V. M. Jiménez-Pérez, J. Magull, A. Ringe and P. G. Jones, *Eur. J. Inorg. Chem.*, 2007, 4919.
- 20 F. A. L. Anet, Top. Curr. Chem., 1974, 45, 169.
- 21 Search of the Cambridge Crystallographic Data Base (October 2014), Al-N range 1.657-2.891 Å (mean 1.956 Å), N-Li range 1.352-2.767 Å (mean 2.077 Å).