Publisher: NPG; Journal: Nature: Nature; Article Type: BCA DOI: 10.1038/nature19779

## Reply to Stojanowski et al.



<sup>1</sup>Leverhulme Centre for Human Evolutionary Studies, Department of Archaeology & Anthropology, University of Cambridge, Fitzwilliam Street, Cambridge CB2 1QH, UK.

In the accompanying Comment<sup>1</sup>, Stojanowski *et al.* challenge the evidence for inter-group conflict at Nataruk<sup>2</sup>. They make two arguments—first, that the lesions in three crania are due to soil compression; second, that there is a correlation between body position and age, reflecting different burial traditions. We believe that their interpretation is incorrect in both counts.

Stojanowski *et al.* argue that the fractures in three Nataruk skulls do not meet forensic criteria, the criteria are unreliable, and the location of lesions is linked to skull/face orientation. We focus on cranium 71264, argued by Stojanowski *et al.* as an example of soil compression effects. This skull has two primary lesions (challenging any site-wide association between skull trauma location and face orientation), one on the left temporoparietal area, another on the frontal bone; these show clear in-bending of bone at impact-point and out-bending of affected regions, radiating fractures, directional change,

<sup>&</sup>lt;sup>2</sup>Turkana Basin Institute, PO box 24467-00502, Nairobi, Kenya.

<sup>&</sup>lt;sup>3</sup>National Museums of Kenya, Jomo Kenyatta House, PO box 152-30500, Lodwar, Kenya.

<sup>&</sup>lt;sup>4</sup>Departamento de Prehistoria y Arqueologia, UNED, c/ Paseo Senda del Rey, 7, 28040 Madrid, Spain.

<sup>&</sup>lt;sup>5</sup>National Museums of Kenya, PO box 40658-00100, Nairobi, Kenya.

<sup>&</sup>lt;sup>6</sup>Jomo Kenyatta University of Agriculture and Technology, PO box 62000-00200, Nairobi, Kenya.

<sup>&</sup>lt;sup>7</sup>Research School of Earth Sciences, The Australian National University, Building 142, Mills Road, Acton, Australian Capital Territory 2601, Australia.

<sup>&</sup>lt;sup>8</sup>Research Laboratory for Archaeology and the History of Art, University of Oxford, Dyson Perrins Building, South Parks Road, Oxford OX1 3QY, UK.

<sup>&</sup>lt;sup>9</sup>Australian Research Centre for Human Evolution, Environmental Futures Research Institute, Griffith University, 170 Kessels Road, Nathan, Queensland 4111, Australia.

<sup>&</sup>lt;sup>10</sup>Department of Geology, Anna University, Chennai, Tamil Nadu 600025, India.

Publisher: NPG; Journal: Nature: Nature; Article Type: BCA DOI: 10.1038/nature19779

depressed adhering bone fragments, lesions across sinuses, evulsion of molars, and differential lateral displacement of maxilla and mandible. We know of no selectively targeted, multi-directional soil compression agent that could account for the observed pattern in this, or other Nataruk crania. The criteria used for identification of perimortem trauma are not only consistent with the literature<sup>3,4</sup>, but clearly identify multiple traumatic lesions at Nataruk<sup>2</sup>.

Stojanowski et al. also argue that prone individuals at Nataruk are older than the rest, and that the site must be a cemetery, with sequential burial traditions, to have preserved the remains. To support this, they (1) claim that there are four prone individuals at Nataruk, not two, by redefining the word 'prone' thus including two skeletons found flexed on their side<sup>2</sup>; (2) dismiss the absence of burial pits or the shells/fish adhering onto the bones as irrelevant; (3) do not account for the fact that bodies and body parts were strewn over 20,000 m<sup>2</sup>; (4) incorrectly state that the majority of human remains at Nataruk were articulated and uncommingled; (5) argue for 'limited variation in body positioning' (figure 1 from ref. 2 shows clearly the diversity of position/direction of bodies, limbs, heads and faces at Nataruk); and (6) draw parallels with cemeteries in Sudan<sup>5</sup>. These are rich multi-stratified, younger sites (~8000–2000 years BP), in one of which earlier burials (with residual burial pits) were prone, later Neolithic ones, flexed<sup>3</sup>. However, prone skeletons in Sudan, although rare, are found across time—from 30,000–20,000 years ago at Wadi Kubbaniya<sup>6</sup> to as recent as 1st millennium BC at Jebel Moya<sup>7,8</sup>, and preserved unburied bodies found before in East Africa<sup>9</sup>. In fact, there are no chronological, stratigraphic or internment parallels between the Omdurman sites and Nataruk, as there are no universal East African pre-Neolithic burial traditions.

Nataruk is not a cemetery, the majority of remains were disarticulated and dispersed; the human remains' multiple cranial and post-cranial traumas are more consistent with perimortem traumatic lesions than soil compression, and critically include embedded projectiles. There is no association between body position and age, or face orientation and location of fractures. A case of inter-group conflict remains the best explanation of what happened at Nataruk.

## Publisher: NPG; Journal: Nature: Nature; Article Type: BCA DOI: 10.1038/nature19779

## References

- <unknown>1 Stojanowski, C. M. et al. Contesting the massacre at Nataruk. Nature (2016).</unknown>
- <jrn>2. Lahr, M. et al. Inter-group violence among early Holocene hunter-gatherers of West Turkana, Kenya. Nature 529, 394–398 (2016). Medline CrossRef
- <jrn>3. Buikstra, J. E. & Ubelaker, D. H. Standards for data collection from human skeletal remains. Am J Hum Biol. 44, 120 (1994).
- <edb>4. Knüsel, C. & Smith, M. J. (Eds). The Handbook of the Bioarchaeology of Human Conflict. (Routledge, 2013).</edb>
- <jrn>5. Salvatori, S., Usai, D. & Zerboni, A. Mesolithic site formation and palaeoenvironment along the White Nile (Central Sudan). *Afr. Archaeol. Rev.* 28, 177–211 (2011). <a href="mailto:CrossRef">CrossRef</a>/jrn>
- <br/>
  <box>6. Wendorf, F. & Schild, R. The Wadi Kubbaniya skeleton: A Late Paleolithic burial from southern Egypt 71–74 (Southern Methodist University Press, 1986).</br>
- <br/><bok>7. Addison, F. Jebel Moya (Oxford University Press, 1949).</br>
- <jrn>8. Brass, M. & Schwenniger, J. L. Jebel Moya (Sudan): new dates from a mortuary complex at the southern Meroitic frontier. Azania 48, 455–472 (2013). Medline CrossRef
- <jrn>9. Leakey, L. S. B. The Naivasha fossil skull and skeleton. Journal of the East African Natural History Society 16, 169–177 (1942).